

NuLYTELY Colonoscopy Prep

Patient	Doctor	-		
Location	Arriva	l Time	AM PM	
Date of Procedure				
make sure that it is ok stop 5 days prior to pi	oumadin, Plavix, Xarelto, Effient, Bril ay to stop them for the recommend ocedure. If you take Xarelto or Eliqui please contact our office. Do not stopedure.	ed number of days is stop 2 days prior	s: If you take Coumadin, r to procedure. If your p	Plavix, Effient or Brilinta rescribing doctor says that
	PROCEDURE: Discontinue eating any mall seeds in it. Also avoid iron table			•
(black or with sugar, be apple juice and clear solid (except red or pur	OCEDURE: Start clear liquid diet all da out no creamers, even non-dairy crea oda (Sprite, 7-up, ginger ale, etc.). Yo ple) and lemonade without pulp. No ure you drink plenty of clear liquids the	mers), tea, clear fo ou may also have (milk or milk produ	ruit juices without pulp s Gatorade, popsicles (exc ucts. This includes SlimFa	such as white grape or ept red or purple), Kool-
4:00 P.M.	Take 2 Bisacodyl tablets or DULCOLAX (over the counter) with water. Select your NuLYTELY flavor of choice and add water to the top line of disposable container. Apply cap and shake until dissolved. Place in refrigerator.			
5:00 P.M.	Drink at least HALF of the solution. Drink one 8-ounce glass every ten minutes			
	OUR PROCEDURE: 6 hours prior to the appleted at least 4 hours prior to your	•	drink the remaining por	tion of the solution. The
DO NOT EAT OR DRI	NK ANYTHING 4 Hours before your	procedure.		
No gum/mints/hard c	andies are allowed on the day of you	r procedure.		
If you take blood pres	sure medications, you may take ther	n the morning of t	he procedure with a sm	all sip of water.

If you have any questions, please call 704-377-4009.

Someone at least 18 years of age or older must bring you to the procedure, remain at the facility during your procedure, and

take you home after the procedure. Otherwise your procedure will be canceled.