

EGD (Upper Endoscopy) Prep

Patient: _____

Doctor: _____

Location: _____

Date of Procedure: _____

Arrival Time: _____

Please Note: These instructions are for patients who are having an EGD (Upper Endoscopy) only. If you are also having a colonoscopy, please follow the colonoscopy prep instructions instead of the instructions below.

- Semaglutide - (ozempic, Wegovy, Rybelsus)
- Dulaglutide - (Trulicity)
- Exenatide Injection - (Bydureon, Byetta)
- Tirzepatide - (Mounjaro, Zepbound)
- Phentermine

7 days

- Coumadin
- Plavix
- Effient
- Brilinta

5 days

- Xarelto
- Eliquis

2 days

- Liraglutide - (Victoza, Saxenda)
- Exenatide Oral - (Bydureon, Byetta)

Stop taking morning of

If you normally take the following medications, please contact the prescribing physician to make sure it is okay to stop them for the recommended number of days.

*If your prescribing doctor says that it is not okay to stop, please contact our office. **Do not stop aspirin for the procedure.**

DO NOT EAT OR DRINK ANYTHING AFTER MIDNIGHT THE EVENING BEFORE YOUR PROCEDURE.

THE MORNING OF YOUR PROCEDURE, you may have small sips of water up until 2 hours before your procedure.

No gum/mints/hard candies are allowed on the day of your procedure.

Blood pressure and heart medications may be taken the morning of your test with a small sip of water.

Someone at least 18 years of age must bring you to the procedure, remain at the facility during your procedure, and take you home after the procedure. Otherwise, your procedure will be canceled.

If you have any questions, please call (704) 377-4009.